



The Transitional Changes Between the Fluvial and Marine Environments as Indicated by Holocene Fauna in Siba at Basrah, Southern Iraq

Hayder Mohammed Jali^{1*} , Abbas Hameed Mohammed ² 

^{1,2} Department of Geology, College of Science, University of Basrah, Basrah, Iraq.

Article information

Received: 13- Jan -2024

Revised: 22- Feb -2024

Accepted: 21- Apr -2024

Available online: 01- Jul – 2025

Keywords:

Recent sediments

Holocene

Foraminifera

Ostracoda

Basrah.

Correspondence:

Name: Hayder Mohammed Jali

Email: haydermrof@gmail.com

ABSTRACT

The study is conducted at one point in Siba City, Basrah to describe its fauna content and texture. The current study attempts to reveal the transitional stage between marine and fluvial environments. Variation in sediment texture provides an index of environmental transformation in the Recent (Holocene Epoch). From grain size analysis, we notice an increase in the percentage of silt and a decrease in the percentage of sand in meters (1-3), a gradual increase in the percentage of sand and a gradual decrease in the percentage of silt in meters (4-6), and in meter 7 there is a high increase in sand and a high decrease in silt, and a gradual decrease in clay from meter 1 to meter 7, a decrease in the percentage of Total Organic Carbon (TOC) is also noted. Recent fauna reinforces sedimentary evidence that marine regression began at depths 5 and 7 m, marine species decrease significantly, and one of the most important marine groups is the Miliolina (Suborder) from foraminifera, where Triloculina, Quinqueloculina, and Spiroloculina are found; on the other hand, describe other groups; Nodosariina, Rotaliida, Textulariina, Polymorphinina and Nodosariina in addition to Ostracoda.

DOI: [10.33899/earth.2024.145028.1184](https://doi.org/10.33899/earth.2024.145028.1184), ©Authors, 2025, College of Science, University of Mosul.

This is an open-access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

التغيرات الانتقالية بين البيئتين النهرية والبحرية بدلالة مستحاثات الهولوسين في السيبة، البصرة، جنوب العراق

حيدر محمد جالي¹  ، عباس حميد محمد²

^{1,2}قسم علوم الأرض، كلية العلوم، جامعة البصرة، البصرة، العراق.

الملخص	معلومات الارشيف
<p>أجريت الدراسة في نقطة واحدة في السيبة، مدينة البصرة لوصف نسيجها ومحاتوتها من المستحاثات. تحاول الدراسة الحالية الكشف عن المرحلة الانتقالية بين البيئتين النهرية والبحرية. توفر الاختلافات الرئيسية في نسيج الرواسب مؤشرًا للتحول البيئي في فترة (الهولوسين) الأخيرة. ومن تحليل حجم الحبيبات نلاحظ زيادة في نسبة الغرين وانخفاض في نسبة الرمل بالأمتار (3-1) وزيادة تدريجية في نسبة الرمل وانخفاض تدريجي في نسبة الغرين بالأمتار (4-6)، وفي المتر 7 هناك زيادة عالية في الرمل وانخفاض كبير في الغرين، وانخفاض تدريجي في الطين من المتر 1 إلى المتر 7، كما يلاحظ انخفاض في نسبة الكربون الكلي العضوي. وما يعزز الأدلة الرسوبيّة المستحاثات الحديثة ويدل على أن التراجع البحري بدأ على أعمق 5 و 7 م، تناقصت الأنواع البحرية بشكل ملحوظ، ومن أهم المجموعات البحرية هي <i>Miliolina</i> (الحدود الفرعية) من المنخربات، حيث تم العثور على <i>Triloculina</i>, <i>Quinqueloculina</i>, and <i>Spiroloculina</i>. ومن ناحية أخرى، تم وصف المجموعات الأخرى؛ <i>Rotaliida</i> و <i>Nodosariina</i> و <i>Textulariina</i> و <i>Polymorphinina</i> و <i>Ostracoda</i>.</p>	<p>تاريخ الاستلام: 13-يناير-2024</p> <p>تاريخ المراجعة: 22-فبراير-2024</p> <p>تاريخ القبول: 21-ابril-2024</p> <p>تاريخ النشر الالكتروني: 01-يوليو-2025</p>
	<p>الكلمات المفتاحية:</p> <p>الرواسب الحديثة</p> <p>الهولوسين</p> <p>فورامينفيرا</p> <p>اوستراكودا</p> <p>البصرة</p>
	<p>المراسلة:</p> <p>الاسم: حيدر محمد جالي</p> <p>Email: haydermrof@gmail.com</p>

DOI: [10.33899/earth.2024.145028.1184](https://doi.org/10.33899/earth.2024.145028.1184), ©Authors, 2025, College of Science, University of Mosul.
This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Sand, silt and clay are components of the Holocene sediments in southern Iraq (Al-Jabberi, 2010; Awadh and Ali, 2011 and Al-Jumaily, 1994). Jassim and Goff (2006) proved that deltaic, aeolian and fluvial sediments are mostly present in the Mesopotamian plain. Holocene sediments split into marine facies and fluvial recent sediments (Al-Jabberi, 2017); the Holocene freshwater deposits of the lower Mesopotamian plain comprise a thin transgressive marine unit extending inland about 250 km from the present coastline (Hammar Formation) (Plaziat and Younis, 2005). Hudson *et al.* (1957) described the Hammar Formation from well No. 13 where there are recent marine sediments up to 6.5 m thick consisting of clay in the upper part and silt and sand in the lower part. This formation can be considered one of the last Quaternary marine transgressions, which are recorded world-wide during the Middle Holocene (Karim *et al.* 1994).

Al-Hemidawy (2020) identified some species of Ostracoda in Al-Delmaj Marsh area. Abdulwahhab (2023) gave systematics for some of foraminifera and Ostracoda in southern Iraq. Al-Furaiji (2022) named some species of foraminifera and Ostracoda in southern Iraq.

There are extensive published and written articles about Holocene deposits in southern Iraq, such as Salman and Al-Mussawy (1994) and Khan *et al.* (1992) referenced the texture of the Holocene deposits as silt and clay where extended from the surface sediments to 18 m depth, while the sand fraction is dominant after 18 m depth. The present study aims to demonstrate the importance of studying fauna assemblages as an important branch in identifying the environments between the fluvial recent sediments and marine sediments at Basrah City.

Study area

The location of the study area represents the Siba area, southern Iraq between latitudes ($30^{\circ}13'04.9''$ - $30^{\circ}20'16.5''$) north and longitudes ($48^{\circ}15'39.0''$ - $48^{\circ}23'03.7''$) east (Fig. 1). South of the city of Basrah, on the Iraqi bank of the Shatt Al-Arab, sedimentation increases, and erosion increases on the Iranian bank. This region consists mainly of alluvial deposits of the Tigris, Euphrates, and Karun rivers and their tributaries.

Materials and methods

The current study includes depths ranging from 1 to 7 meters at one point. From the core, (33) cutting samples are selected according to the differences in the textures, colors of the beds, and grain size analysis following Folk (1974) and Total Organic Carbon (TOC%) is determined according to methods of Nelson and Sommers (1996). After that, the samples were left in water for three days to soak, then washed under the tap water in 230 mesh sieve to picking the fauna, and they were stored in micropaleontological slide, two groups for this study are identified (foraminifera and ostracod).



Fig. 1. The study area map, Iraq, Basrah City.

Results

Sediments Texture analysis

The study area sediments are dated back to the Holocene. Through the current study, because it is a part of the fresh water of Shatt Al-Arab, they are characterized essentially by sandy silt and clayey silt respectively according to the texture classification of Folk (1974).

After leaving the marshes at Al-Qurna area, Tigris and Euphrates Rivers joined each other to create the river of Shatt Al-Arab, which flows in somewhat winding channels. Hor Al-Hammar feeds Shatt Al-Arab through the Garmat Ali channels north of Basrah (Yacoub, 2011).

The input of the Karun River affects Shatt Al-Arab; as a result of the junction that occurred north of Siba, there is a sudden increase in the amount of mud that was suspended in the water of the Shatt Al-Arab. The tides affect the level of Shatt Al-Arab, principally downstream of Basrah City (Yacoub, 2011). Fine sediments textures are characteristic of tidal flats, sometimes sand and mostly clay and silt. Table (1) summarizes the textural results; it is found that silty sand and clayey silt are the dominant textures.

Table 1: Grain size analysis and TOC% of cutting sediments in Siba area.

NO.	Depth (cm)	Sand%	Silt %	Clay%	TOC%
1	(1m) 4-9	2	75	23	0.79
2	(1m) 22-25	1	81	18	1.05
3	(1m) 37-41	12	70	18	1.15
4	(1m) 49-52	2	78	20	0.55
5	(1m) 63-66	2	88	10	0.46
6	(2m) 127-130	3	82	15	0.55
7	(2m) 145-150	5	77	18	0.45
8	(2m) 166-169	3	62	35	0.86
9	(2m) 187-190	2	83	15	0.49
10	(3m) 200-205	13	79	8	0.70
11	(3m) 219-223	4	91	5	0.63
12	(3m) 245-249	12	78	10	0.46
13	(3m) 272-275	9	81	10	0.21
14	(3m) 292-962	5	77	18	0.58
15	(4m) 300-304	9	76	15	0.57
16	(4m) 327-331	11	81	8	0.57
17	(4m) 344-348	87	5	8	0.39
18	(4m) 370-374	40	42	18	0.40
19	(4m) 393-397	74	11	15	0.37
20	(5m) 400-403	17	68	15	0.37
21	(5m) 423-426	40	48	12	0.36
22	(5m) 452-455	37	48	15	0.29
23	(5m) 472-476	41	49	10	0.23
24	(5m) 495-498	25	62	13	0.39
25	(6m) 510-514	53	42	5	0.27
26	(6m) 534-538	36	54	10	0.33
27	(6m) 555-559	29	63	8	0.44
28	(6m) 572-576	31	61	8	0.42
29	(6m) 590-594	44	46	10	0.41
30	(7m) 600-604	61	31	8	0.25
31	(7m) 619-623	75	15	10	0.43
32	(7m) 638-642	40	52	8	0.41
33	(7m) 669-673	81	16	3	0.21

Recent Faunal Distribution

1. Foraminifera

Foraminifera are among the important groups identified in this area due to its wide environmental range from bottom of sea to continental regions (Boltovskoy and Wright, 1976). Depending on the classification of Loeblich and Tappan (1964), many species are recorded and identified (26 species) belonging to 13 genus as illustrated in Table (2) and Plates (1,2,3,4 and 5). These species are: *Ammonia tepida*, *A. beccarii*, *A. dentata*, *Asterorotalia inflata*, *Brizalina striatula*, *Caribeanella polystoma*, *Elphidium advenum*, *E. crispum*, *E. discoidale*, *E. hispidilum*, *E. incertum*, *E. lessonii*, *Fissurina marginata*, *Lagena perlucida*, *Lagena striata*, *Nonion elongatum*, *Parrellina hispidula*, *Quinqueloculina agglutianans*, *Q. angulta*, *Q. parvula*, *Q. polygona*, *Q. seminulum*, *Q. stelligera*, *Triloculina oblonga*, *S. exima*, *Spirosigmoilina tenuis*. Near shore or lagoon that are shallow when depth 1-20 m live *Ammonia beccarii*, *Elphidium incertum* (recent species) (Murray, 1969).

Salinity is the wide range reflected by fauna assemblages, carbonate ions in highly saline water are high, 40% and more is the percentage of salinity, appear to favor the porcelaneous *Miliolina* (especially the Miliolidae and Nubecularidae, e.g., *Triloculina*, *Quinqueloculina*).

Table 2: Distribution of Foraminifera species in the 7 meters studied section in Siba (the empty square is fauna-free).

Depth (cm)	<i>Triloculina oblonga</i>	<i>Spirosgmoinitina tenuis</i>	<i>Spiroloculina exima</i>	<i>Q.stelligera</i>	<i>Q.seminulum</i>	<i>Q.polygona</i>	<i>Q.parvula</i>	<i>Q.angulata</i>	<i>Quinqueloculina agglutinans</i>	<i>Parcellina hispidula</i>	<i>Nonion elongatum</i>	<i>Lagenia striata</i>	<i>Lagenia perlucida</i>	<i>Fissurina marginata</i>	<i>E.lessonii</i>	<i>E.incertum</i>	<i>E.hispidulum</i>	<i>E.edicoidale</i>	<i>E.crispum</i>	<i>Ephistium advenum</i>	<i>Caribeanelleta polystoma</i>	<i>Brizalina striatula</i>	<i>Asterorotalia inflata</i>	<i>A.lepida</i>	<i>A.dentat</i>	<i>Ammonia beccarii</i>		
(1m) 4-9	*	*	*																									
(1m) 22-25	*	*	*																									
(1m) 37-41																												
(1m) 49-52																												
(1m) 63-66																												
(2m) 127-130	*																											
(2m) 145-150	*	*																										
(2m) 166-169	*																											
(2m) 187-190	*	*	*	*																								
(3m) 200-205	*		*	*																								
(3m) 219-223																												
(3m) 245-249	*	*	*	*																								
(3m) 272-275	*		*																									
(3m) 292-962	*		*																									
(4m) 300-304	*		*																									
(4m) 327-331	*		*																									
(4m) 344-348	*																											
(4m) 370-374	*	*	*	*																								
(4m) 393-397	*	*	*	*																								
(5m) 400-403	*	*	*																									
(5m) 423-426	*	*	*																									
(5m) 452-455	*	*	*																									
(5m) 472-476	*		*																									
(5m) 495-498	*	*	*																									
(6m) 510-514	*		*	*	*	*	*	*																				
(6m) 534-538	*	*	*	*	*																							
(6m) 555-559	*	*	*																									
(6m) 572-576	*	*	*		*																							
(6m) 590-594	*	*	*		*																							
(7m) 600-604	*	*	*	*	*	*	*	*																				
(7m) 619-623	*	*	*																									
(7m) 638-642	*	*	*																									
(7m) 669-673	*	*	*																									

2. Ostracoda

In this research, the second group was the Ostracods, the diagnosed of Ostracoda species is depending on Moore and Pitrat, (1961). 12 species belonging to 9 genus were identify (Table 3 and Pl. 6 and 7), these species are: *Alocopocythere reticulata*, *Candoniella albicans*, *Candoniella simpsoni*, *Carinocythereis indica*, *Cyprideis torosa*, *Cyprideis var. torosa*, *Darwinula cylindrical*, *Hemicytheridea paiki*, *Hemicytheridea reticulata*, *Leguminocythereis*

papuensis, *Neomonoceratina delicata*, *Phlyctenophora orientalis*. The species were represented a marine taxon Shareef and Mahdi, (2015) and Moore and Pitrat, (1961).

Table 3: Distribution of Ostracoda species on the 7 meters studied section in Siba (the empty square is valid fauna-free).

<i>Depth (cm)</i>	<i>Phlyctenophora orientalis</i>	<i>Neomonoceratina delicata</i>	<i>Leguminocythereis papuensis</i>	<i>Hemicytheridea reticulata</i>	<i>Hemicytheridea palki</i>	<i>Darwinula cylindrical</i>	<i>Cyprideis</i> var. <i>torosa</i>	<i>Carinocythereis indica</i>	<i>Candoniella simpsoni</i>	<i>Candoniella albicans</i>	<i>Alocopocythere reticulata</i>
(1m) 4-9											
(1m) 22-25											
(1m) 37-41											
(1m) 49-52											
(1m) 63-66											
(2m) 127-130											
(2m) 145-150											
(2m) 166-169											
(2m) 187-190											
(3m) 200-205											
(3m) 219-223											
(3m) 245-249	*										
(3m) 272-275							*				
(3m) 292-962							*				
(4m) 300-304							*				
(4m) 327-331							*				
(4m) 344-348	*						*				
(4m) 370-374							*				
(4m) 393-397							*				
(5m) 400-403		*					*				
(5m) 423-426	*						*				
(5m) 452-455	*						*				
(5m) 472-476							*				
(5m) 495-498							*				
(6m) 510-514				*			*				
(6m) 534-538				*			*				
(6m) 555-559				*			*				
(6m) 572-576			*	*			*				
(6m) 590-594		*		*			*				
(7m) 600-604		*		*			*				
(7m) 619-623		*		*			*				
(7m) 638-642			*	*			*				
(7m) 669-673			*	*			*				*

Table 4: Systematic of foraminifera.

Kingdom	Phylum	Order	Family	Genus
Chromista Cavalier-Smith, 1981	Foraminifera d Orbigny, 1826	Rotaliida Delage and Herouard, 1896	Ammoniida Saidova, 1981	<i>Ammonia</i> Brunnich, 1771
			Elphidiidae Galloway, 1933	<i>Asterorotalia</i> Hofker, 1950
			Notorotaliidae Hornbrook, 1961	<i>Elphidium</i> Montfort, 1808
			Bolivinidae Cushman, 1927	<i>Parrellina</i> Thalmann, 1951
			Planorbulinidae Schwager, 1877	<i>Brizalina</i> Costa, 1856
			Nonionidae Schultze, 1854	<i>Caribeanella</i> Bermudez, 1952
			Ellipsolagenidae A. Silvestri, 1923	<i>Nonion</i> Montfort, 1808
		Polymorphinina Mikhailevich, 1980	Lagenidae Reuss, 1862	<i>Fissurina</i> Reuss, 1850
		Nodosariina Calkins, 1926	Hauerinidae Schwager, 1876	<i>Lagen</i> Walker & Jacob, 1798
		Miliolina Williamson, 1858		<i>Quinqueloculina</i> d'Orbigny, 1826
				<i>Triloculina</i> d'Orbigny, 1826
				<i>Spirosigmoilina</i> Parr, 1942
				<i>Spiroloculinidae</i> Wiesner, 1920
				<i>Spiroloculina</i> d'Orbigny, 1826

Table 5. Systematic of Ostracoda.

Kingdom	Phylum	Class	Family	Genus
Animalia Linnaeus, 1785	Arthropoda Siebold and Stannius, 1845	Ostracoda Latreille, 1806	Trachyleberididae Sylvester-Bradley, 1948	<i>Carinocythereis Ruggieri</i> , 1955. <i>Leguminocythereis Howe</i> , 1936. <i>Keijella Ruggieri</i> , 1967. <i>Alocopocythere Siddiqui</i> , 1971.
			Cytherideidae Sara, 1925	<i>Cyprideis Jones</i> , 1857.
			Cytheridae Baird, 1850	<i>Neomonoceretina Kingma</i> , 1948.
			Hemicytheridae Puri, 1953	<i>Hemicytheridea Kingma</i> , 1948.
			Candonidae Kaufmann, 1900	<i>Phlyctenophora Brady</i> , 1880.
			Darwinulidae Brady & Robertson, 1885	<i>Candoniella Schneider</i> , 1956. <i>Darwinula Brady & Robertson</i> , 1885.

Discussion

The present study aims to recognize assemblages of fauna during the sedimentation of the Holocene in Basrah City. From grain size analyses in 1m to 4m depth, clay and silt fractions are predominant with clayey silt texture, while sand fraction started to increase from 42% at a depth interval of 4 m to 65 % at 7m, with silty sand texture (Table 1).

The organisms (Ostracoda and Foraminifera) are examined. These are the assemblages of Recent fauna that have been identified and relied upon, three stages are distinguished in this research from bottom to top respectively (Tables 2, 3). Interval depth 7 to 6 m represents the first stage, the facies are silty sand, which resembles the facies of the coastal marine environment. The similarity of fauna also may reflect a combination of sedimentation. Interval depths 6 to 3 m represent the second stage, and facies was (sandy silt), saline water is what assemblages' fauna indicated to like *Quinqueloculina seminulum*, *Hemicytheridea reticulata*, *Triloculina oblonga*, *Carinocythereis indica*. etc. Interval depth 3 to 0 m represents the third stage, which represents the decline of seawater and the advance of river water accompanied with a change in the existing fauna.

Conclusions

1- Many foraminifera and ostracoda species are identified that are indicators for regression.

2- Three sedimentary textures are distinguished in the study area (muddy silt, sandy silt and silty sand).

3- The presence of the species *Ammonia beccari* at most depths indicates the mixed environment in the study area as it refers to the shallow marine environments to the brackish environment, and it is adapted to living in those environments.

4- Marine regression started at 5 m in Siba.

References

Abdulwahhab Z.M., 2023. Depositional and Stratigraphic Study of Mauddud Formation in Selected Oil Fields, Southern Iraq. PhD Thesis, Basrah University, College of Science, 138 P.

Al-Hemidawy F.N.M., 2020. Recent Fauna (Ostracoda, Foraminifera and Mollusca) and Geochemical of Al-Delmaj Marsh Area, Iraq. PhD Thesis, Basrah University, College of Science, 212 P.

Al-Furaiji S.A.M., 2022. A Paleontological and Sequence Stratigraphy Study of Yamama Formation in Selected Wells Southern Iraq. Ph.D. Thesis, Basrah University, College of Science, pp.212.

Al-Jabberi, M.H.A., 2010. Sedimentological and Environmental Aspects of Subsurface Basrah Sediments-South Iraq. Mesopotamian Journal of Marine Science, 25(2), pp. 176-187.<https://doi.org/10.58629/mjms.v25i2.203>

Al-Jaberi, M.H., 2017. Clay Minerals Variations in Quaternary Sediments of Basrah City-Iraq. Journal of Earth Science Research-Published by Canadian Center of Science and Education, 6(2), pp. 41-55. <https://doi.org/10.5539/esr.v6n2p41>

Al-Jumaily, W.A., 1994. Quaternary Ostracods in Southern Iraq. Unpublished PhD Thesis, Baghdad University, 212 P.

Awadh, S.M., Ali, M.O. and Ali, R.A., 2011. Mineralogy and Palynology of the Mesopotamian Plain Sediments, Central Iraq. Arabian Journal of Geosciences, 4, pp. 1261-1271. <https://doi.org/10.1007/s12517-010-0161-y>.

Boltovskoy, E., and Wright, R., 1976. Recent Foraminifera. Springer Science Business Media Dordrecht, 515 P. <http://dx.doi.org/10.1007/978-94-017-2860-7>.

Hudson, R.G.S., Eames, F.E., and Wilkins, G.L., 1957. The Fauna of Some Recent Marine Deposits Near Basrah- Iraqi Geological Journal, 94(5), pp. 393-401. <https://doi.org/10.1017/S0016756800069429>

Jassim, S.Z., and Goff, J.C., 2006. Geology of Iraq. Dolin, distributed by Geological Society of London, 341 P. <http://dx.doi.org/10.4236/oje.2014.411060>

Karim, S.A., Al-Jumaily, W.A., and Al-Sheikhly, S.S., 1994. Quaternary Foraminifera of Basrah Area. Iraqi Geological Journal, 27(2), pp. 1-13.

Khan, N., Albadran, A.A., and Albadran, B., 1992. Some Engineering Aspects of the Alluvial Deposits at Basrah Region. Iraqi Geological Journal, 25(1), pp. 80-91.

Loeblich, A.R. and Tappan, H., 1964. Thecamoebians and Foraminifera. Kansas Press Lawrence, Kans, 900 P.

Moore, R.S. and Pitrat, C.W. 1961. Treatise, on Invertebrate, Paleontology, Arthropoda, (Crustacea Ostracoda), 442 P., 334 figs. Geol. Soc. Amer. and Univ. of Kiel, fig. 334, 442 pp.

Murray, J.W., 1969. Recent Foraminifers From the Atlantic Continental Shelf of the United States. Micropaleontology, 15, pp. 401-419.

Nelson, D.W. and Sommers, L.E., 1996. Total Carbon, Organic Carbon, and Organic Matter. In: Sparks, D.L., et al., Eds., Methods of Soil Analysis. Part 3. Chemical Methods, SSSA Book Series No. 5, SSSA and ASA, Madison, WI, pp. 961-1010.

Plaziat, J.C. and Younis, R.W., 2005. The Modern Environments of Molluscs in Southern Mesopotamia, Iraq: A guide to paleogeographical reconstructions of Quaternary fluvial, Palustrine and Marine Deposits. Carnets de Geologie, 15, pp. 1-18. <http://dx.doi.org/10.4267/2042/1453>

Salman, H. and Al-Mussawy, S., 1994. Some Sedimentological Aspects of the Shatt Al-Arab Levee Sediments. Iraqi Geological Journal, 27(3), pp. 20-39.

Shareef, N.F. and Mahdi, M.M., 2015. Studying of Recent Environments in Faw, Khor Al-Zubair and Um- Qaser Areas, Southwestern Arabian Gulf, Basrah, Iraq. Journal of Basrah Researches, 41(2), pp. 1-14.

Folk, R.L. 1974. Petrology of Sedimentary Rocks, Hemphill Publishing Company, Austin, Texas, 183 P. <http://dx.doi.org/10.4236/ijg.2015.63019>

Yacoub, S.Y. 2011. Stratigraphy of the Lower Mesopotamia Plain. Iraqi Bull. Geol. Min. Special Issue, No.4, Geology of the Mesopotamia Plain, pp. 47-82. <http://dx.doi.org/10.4236/ad.2015.31003>

Plate 1Scale 1bar = 200 μ m.

- 1- *Ammonia beccarii* Linnaeus, 1758. 40x, side view.
- 2- *Ammonia tepida* Cushman, 1926. 40x, side view.
- 3- *Ammonia dentata* Parker and Jones. 40x, side view.
- 4- *Asterorotalia inflata* Millett, 1904. 40x, side view.
- 5- *Brizalina striatula* Cushman. 40x, side view.
- 6- *Caribbeanella polystoma* Bermúdez, 1952. 40x, side view.







Plate 1.

Plate 2

Scale 1bar = 200 μ m.

- 1- *Elphidium advenum* Cushman, 1922. 40x, side view.
- 2- *Elphidium crispum* Linnaeus, 1758. 40x, side view.
- 3- *Elphidium hispidulum* Cushman, 1936. 40x, side view.
- 4- *Elphidium incertum* Williamson, 1858. 40x, side view.
- 5- *Elphidium lessonii* d'Orbigny, 1839. 40x, side view.
- 6- *Elphidium discoidale* d'Orbigny, 1839. 40x, side view.





Plate 2**1****2****3****4****5****6**

Plate 3

Scale 1bar = 200 μ m.

- 1- *Parcellina hispidula* Cushman, 1936. 40x, side view.
- 2- *Nonion elongatum* d'Orbigny, 1852. 40x, side view.
- 3- *Lagena striata* d'Orbigny, 1839a. 40x, side view.
- 4- *Lagena perlucida* Montagu, 1803. 40x, side view.
- 5- *Fissurina marginata* Montagu, 1803. 40x, side view.
- 6- *Quinqueloculina agglutinans* d'Orbigny, 1839. 40x, side view.






Plate 3

Plate 4

Scale 1bar = 200 μ m.

- 1- *Quinqueloculina polygona* d'Orbigny, 1839. 40x, side view.
- 2- *Quinqueloculina parvula* Schlumberger, 1894 40x, side view.
- 3- *Quinqueloculina angulata* Williamson, 1858. 40x, side view.
- 4- *Quinqueloculina stelligera* Schlumberger, 1893. 40x, side view.
- 5- *Quinqueloculina seminulum* Linne, 1758. 40x, side view.
- 6- *Triloculina oblonga* Montagu, 1803. 40x, side view.




Plate 4

Plate 5: Scale 1bar = 200 μm . (1) *Spiroloculina exima* Cushman, 1922. 40x, side view. (2) *Spirosigmoilina tenuis* Cžžek, 1848. 40x, side view.

Plate 5




Plate 6

Scale 1bar = 200 μ m.

- 1- *Alocopocythere reticulata* Hartmann, 1964. 40x, side view.
- 2- *Carinocythereis indica* Jain, 1978. 40x, side view.
- 3- *Candoniella albicans* Sharpe, 1897. 40x, side view.
- 4- *Leguminocythereis papuensis* (Brady) Jain, 1978. 40x, side view.
- 5- *Cyprideis torosa* Jones, 1850. 40x, side view.
- 6- *Cyprideis Var torosa* Jones 1850. 40x, side view.






Plate 6

Plate 7Scale 1bar = 200 μ m.

- 1- *Neomonoceratina delicate* Ischizaki and Kato, 1976. 40x, side view.
- 2- *Hemicytheridea reticulata* Kingma, 1948. 40x, side view.
- 3- *Hemicytheridea paiki* Jain, 1978. 40x, side view.
- 4- *Phlyctenophora orientalis* Brady, 1868. 40x, side view.
- 5- *Candoniella simpsoni* Sharpe, 1897. 40x, side view.
- 6- *Darwinula cylindrica* Straub, 1952. 40x, side view.

Plate 7